



typically consists of commercial monkey biscuits or monkey chow, with limited portions of seasonal fruits (Takenaka *et al.* 2000). Differences in diet between captive and wild environments reportedly influence changes in the composition of LTMs' gut microbiota (Boonkusol *et al.* 2020 Frankel *et al.* 2019; Grant *et al.* 2019; Wills *et al.* 2022).

The diversity of gut microbiota is influenced by a variety of factors, including intrinsic elements such as the age and sex of host species, as well as extrinsic factors like environment and diets (Ying *et al.* 2022). Diet, in particular, plays a crucial role in shaping gut microbiota diversity, as demonstrated by significant alterations in gut microbiota diversity within just a year of transferring LTMs from their natural habitat to captivity (Sawaswong *et al.* 2023). In primates, the gut microbiota comprises bacteria, fungi, protozoa, viruses, and archaea. These microorganisms play essential roles in metabolizing complex nutrients (Chen *et al.* 2017), maintaining immune system function (Muegge *et al.* 2011), and regulating hormone activity (Martin *et al.* 2019). Additionally, the gut microbiota serves as a secondary neural network, facilitating bidirectional communication between the nervous system and gut microbiota, known as the Microbiota-Gut-Brain-Axis (MGBA) (Carabotti *et al.* 2015). This communication occurs through various mechanisms, including the stimulation of cytokine expression, production of microbial metabolites such as short-chain fatty acids (SCFAs), and tryptophan metabolism (Carbia *et al.* 2023). Moreover, the gut microbiota contributes to the synthesis of neurotransmitters like serotonin which influences gastrointestinal motility and homeostasis (Yano *et al.* 2015), and  $\gamma$ -aminobutyric acid (GABA) which modulates blood pressure and immune function (Pokusaeva *et al.* 2017).

Maintaining the balance of bacteria in the intestine is crucial for healthy bodily function. Dysbiosis, characterized by an imbalance in gut microbiota composition, can lead to the development of idiopathic chronic diarrhea (ICD) in LTMs (Koo *et al.* 2020), immune imbalance in rhesus macaques (Li *et al.* 2020), and has been associated with depressive-like behavior in LTMs serving as a model animal (Wu *et al.* 2022). Consequently, understanding the gut microbiota composition becomes pivotal in assessing LTM health. Despite research efforts focusing on various aspects of LTMs on Tinjil Island, including population dynamics (Perwitasari-Farajallah *et al.* 2023a), behavior (Hasanah *et al.* 2022), and feeding ecology (Perwitasari-Farajallah *et al.* 2023b), studies on gut microbiota composition of semi-wild LTMs have not yet been conducted. Hence, the aim of this study was to assess intestinal bacteria from LTMs on Tinjil Island by isolating and characterizing gut bacteria from their fecal samples using a culture-dependent method. This research provides preliminary insight into the diversity of gut bacteria in semi-wild LTMs from Tinjil Island.

## 2. Materials and Methods

### 2.1. Fecal Sample Collection

In July 2022, we collected a single fecal sample, noninvasively, from four semi-wild LTMs on Tinjil Island, Banten, Primate Research Center, IPB University (Figure 1), with approval from the IPB University Animal Care and Use Committee of Ethics (ACUC-IPB University; no. IPB PRC-19-A012). A sterile wooden stick was used to collect freshly expelled fecal samples, which were then placed into Falcon tubes and refrigerated to preserve their condition for subsequent analysis. Samples were collected from the inner part of the feces, ensuring that they were not exposed to the ground or air.




Figure 1. Fecal sampling locations of the long-tailed macaques on Tinjil Island. 1) Map of Tinjil Island (modified from Perwitasari-Farajallah *et al.* 2023a); 2) Location of fecal sampling points at Base camp and Cage 7; 3) Four fecal samples, one each from four long-tailed macaques: RB1 (Right Basecamp Sample), LB2 and LB3 (Left Base camp Sample), C72 (Cage 7 Sample).

## 2.2. Total Plate Count (TPC)

The TPC method was employed to estimate the population of microorganisms within the growth media by enumerating each bacterial colony. One gram of each LTM fecal sample was homogenized in 9 mL of 0.85% NaCl solution, followed by serial dilution eight times. Subsequently, 0.1 mL of the diluent from  $10^{-4}$  to  $10^{-8}$  serial dilution was spread onto nutrient agar (NA) medium and then incubated aerobically at room temperature ( $25 \pm 2^\circ\text{C}$  for 24-48 hours, with modifications adapted from (Prats *et al.* 2008). This procedure was used to quantify bacterial colonies on each agar plate using TPC, resulting in colony-forming units (CFU/g). Enumeration of the bacterial colonies on each plate was exclusively conducted at dilutions where the total colony count ranged from

30 to 300 bacterial colonies (Zuberer 1994). Colonies numbering less than 30 were classified as “too few to count” (TFTC), while those exceeding 300 were considered “too numerous to count” (TNTC).

## 2.3. Bacterial Isolates Purification

We observed the bacterial colonies on each sample, observing variations in morphology, including form, margin, texture, color, transparency, and elevation, before isolating purified bacteria. We then isolated each colony with distinctive features using the quadrants streak method into the NA plate medium to attain a pure single bacterial colony. The isolates were placed into a refrigerator for further bacteria stain after incubation for about 24-48 hours at  $25 \pm 2^\circ\text{C}$ .

## 2.4. Gram Stain and Biochemical Test

We classified bacteria into two groups using the Gram stain method: Gram-positive and Gram-negative bacteria. Gram-positive bacteria exhibit a purple or blue color when viewed under the microscope, while Gram-negative bacteria appear red (Smith and Hussey 2020). We considered the different bacteria cells' shapes and Gram stain status to move on to the specific biochemical test pathway.

We used biochemical tests to discern various bacteria based on their reactions with different biochemical compounds. Each isolate was identified through a series of biochemical tests following Bergey's Manual of Determinative Bacteriology (Holt *et al.* 1994). Specific biochemical test flowcharts were employed for each bacterial isolate depending on the type of bacteria revealed by Gram staining the shape of the cell. In this study, we used two biochemical test flowcharts: The Gram-positive cocci flowchart and the Gram-positive bacilli catalase flowchart (Supplementary Material Figure S1).

### 2.4.1. Catalase Test for Gram-positive cocci and Gram-positive bacilli

We conducted the slide or drop catalase test to detect the presence of the enzyme catalase in bacteria (Reiner 2013). A positive result was immediately observed when bubbles formed on the slide. Positive results from the catalase test in the Gram-positive cocci group proceeded to the mannitol fermentation test stage (Hanson 2008). Conversely, positive results from the catalase test in the Gram-positive bacilli group led to the starch hydrolysis test (Hussey 2008) with modifications. Colonies resistant to acid decolorization during stain procedures prompted further testing using the acid-fast stain protocol (Hussey and Zayaitz 2013) employing carbolfuchsin as the primary stain, methylene blue as the counterstain, and acid-alcohol as the decolorizing solvent. The

Voges-Proskauer (VP) test determined the organism's ability to produce acetylmethylcarbinol from glucose fermentation (following Mcdevitt's protocol; Mcdevitt 2009) by preparing Methyl Red-Voges Proskauer (MRVP) broth as per the manufacturer's instructions. For the salt tolerance test, we utilized nutrient broth (NB) supplemented with sodium chloride to achieve a salt concentration of 6.5% (6.5 g NaCl per 100 mL NB media) (Bruins *et al.* 2007). Inoculum from a pure culture (aged 18-24 hours) was transferred to sterile 6.5% NaCl broth and incubated at 35-37°C for 24 hours. The presence of turbidity indicated a positive test result.

## 2.5. Hemolysis for all the isolates

We conducted the hemolysis test by inoculating a colony from a fresh (16-18 hours), pure culture onto blood agar plates (BAP) and subsequently incubating them at 35±2°C for approximately 18-24 hours. Bacterial hemolytic reactions are categorized into three types: Beta hemolysis ( $\beta$ ), alpha hemolysis ( $\alpha$ ), and gamma hemolysis ( $\gamma$ ) (Buxton 2016). We conducted our observation with the light shining from behind the plate to interpret the hemolytic reaction of each bacterial streak accurately. Beta hemolysis manifests as a complete lysis of red blood cells, appearing as a clear zone or transparency surrounding the colony. A green or brown discoloration in the medium characterizes alpha hemolysis. Finally, gamma hemolysis is indicated by the absence of a clear zone surrounding the colony on the medium.

## 3. Results

### 3.1. Total Viable Bacteria in LTMs Fecal Samples

According to the total plate count method, the average of viable bacteria found in the fecal samples from the four LTMs was  $1.86 \times 10^9$  CFU/g (Supplementary Material Table S1). However, each sample exhibited varying quantities of bacterial



colonies. Sample C72 displayed the highest number of viable bacteria, whereas Sample RB1 exhibited the lowest.

### 3.2. Bacterial Colony Morphology

Various morphological characteristics were observed among different bacterial colonies grown in each sample (Supplementary Material Table S2, Figure 2). Sample LB3 exhibited the highest number of bacterial isolates, totaling six isolates, while sample C72 had the fewest, with only three isolates. Among the three bacterial isolates from sample C72, only the color differed, while other characteristics remained the same. Additionally, two bacterial isolates with identical colony morphology were identified in sample C72, suggesting they were the same type of bacteria. The colony morphology of bacterial isolates

from samples RB1 and RB2 varied significantly, indicating the presence of different types within these samples. Similarly, sample LB2 contained six isolates, two of which shared similar macroscopic characteristics, namely isolates LB3.1 and LB3.4. Furthermore, isolates with identical morphological characteristics were found across samples, such as isolate RB1.4 with isolate LB3.3 and isolate RB1.1 with isolate LB2.3.

### 3.3. Bacterial Cell Morphology

The characteristics of bacterial cell morphology, including shape, cell arrangement, and Gram group, were assessed (Supplementary Material Table S3, Figure 3). All bacterial isolates were identified as Gram-positive bacteria, predominantly




Figure 2 Bacterial colonies isolated from fecal samples of wild long-tailed macaques on Tinjil Island (1) C72.1 (Cage 7 Sample 2 Isolate 1), (2) C72.2 (Cage 7 Sample 2 Isolate 2), (3) C72.3 (Cage 7 Sample 2 Isolate 3), (4) LB2.1 (Left Base camp Sample 2 Isolate 1), (5) LB2.2 (Left Base camp Sample 2 Isolate 2), (6) LB2.3 (Left Base camp Sample 2 Isolate 3), (7) LB2.4 (Left Base camp Sample 2 Isolate 4), (8) LB2.5 (Left Base camp Sample 2 Isolate 5), (9) LB3.1 (Left Base camp Sample 3 Isolate 1), (10) LB3.2 (Left Base camp Sample 3 Isolate 2), (11) LB3.3 (Left Base camp Sample 3 Isolate 3), (12) LB3.4 (Left Base camp Sample 3 Isolate 4), (13) LB3.5 (Left Base camp Sample 3 Isolate 5), (14) LB3.6 (Left Base camp Sample 3 Isolate 6), (15) RB1.1 (Right Base camp Sample 1 Isolate 1), (16) RB1.2 (Right Base camp Sample 1 Isolate 2), (17) RB1.3 (Right Base camp Sample 1 Isolate 3), (18) RB1.4 (Right Base camp Sample 1 Isolate 4), (19) RB1.5 (Right Base camp Sample 1 Isolate 5). White line on under leftside of every part of figure is a scale.




Figure 3 Positive-gram bacteria isolated from fecal samples of wild long-tailed macaques on Tinjil Island (1) C72.1 (Cage 7 Sample 2 Isolate 1), (2) C72.2 (Cage 7 Sample 2 Isolate 2), (3) C72.3 (Cage 7 Sample 2 Isolate 3), (4) LB2.1 (Left Base camp Sample 2 Isolate 1), (5) LB2.2 (Left Base camp Sample 2 Isolate 2), (6) LB2.3 (Left Base camp Sample 2 Isolate 3), (7) LB2.4 (Left Base camp Sample 2 Isolate 4), (8) LB2.5 (Left Base camp Sample 2 Isolate 5), (9) LB3.1 (Left Base camp Sample 3 Isolate 1), (10) LB3.2 (Left Base camp Sample 3 Isolate 2), (11) LB3.3 (Left Base camp Sample 3 Isolate 3), (12) LB3.4 (Left Base camp Sample 3 Isolate 4), (13) LB3.5 (Left Base camp Sample 3 Isolate 5), (14) LB3.6 (Left Base camp Sample 3 Isolate 6), (15) RB1.1 (Right Base camp Sample 1 Isolate 1), (16) RB1.2 (Right Base camp Sample 1 Isolate 2), (17) RB1.3 (Right Base camp Sample 1 Isolate 3), (18) RB1.4 (Right Base camp Sample 1 Isolate 4), (19) RB1.5 (Right Base camp Sample 1 Isolate 5).

cocci-shaped, with a smaller proportion being bacilli-shaped. Isolate RB1.4 and Isolate LB3.3 exhibited coccobacilli-shaped morphology, intermediate between cocci and bacilli shapes. The arrangement of the bacterial cells varied within each group of Gram-positive cocci and Gram-positive bacilli, with dominance determined through microscopic observation.

### 3.4. Bacterial Identification with Biochemical Tests

Biochemical tests were conducted for the identification of Gram-positive cocci and Gram-positive bacilli, as detailed in Tables 1 and 2, respectively. Based on biochemical test results, only two groups of putative bacterial genera were identified: *Staphylococcus* and *Micrococcus* for the Gram-positive cocci group (Table 1). Isolates C72.2

and C72.3 were identified as *Staphylococcus aureus*, a conclusion supported by their macroscopic colony characteristics (Supplementary Material Table S2). Four bacterial isolates that could not ferment mannitol (indicated by the green OF media color) but exhibited yellow colonies were suspected to be *Micrococcus luteus* bacteria (Supplementary Material Table S4).

Overall, bacterial isolates suspected to be *Staphylococcus* spp. (excluding *Staphylococcus aureus*) shared similar biochemical test characteristics. Similarly, bacterial isolates presumed to be *Micrococcus luteus* species displayed a consistent inability to ferment mannitol. However, a notable difference was observed in colony pigmentation, with suspected *Micrococcus luteus* isolates exhibiting yellow-pigmented colonies while those of suspected *Staphylococcus* spp. (except *Staphylococcus aureus*) tended to have white colonies (Supplementary Material Table S4).

Tabel 1. Biochemical tests on bacterial isolates of Gram-positive cocci

No	Isolate Code	Biochemical Test				Putative species
		Ca	M	Y	G	
1	C72.2	+	+	+	x	<i>Staphylococcus aureus</i>
2	C72.3	+	+	+	x	<i>Staphylococcus aureus</i>
3	LB2.2	+	-	+	-	<i>Micrococcus luteus</i>
4	LB2.3	+	-	-	x	<i>Staphylococcus sp.</i>
5	LB3.2	+	-	+	-	<i>Micrococcus luteus</i>
6	LB3.5	+	-	-	x	<i>Staphylococcus sp.</i>
7	LB3.6	+	-	+	-	<i>Micrococcus luteus</i>
8	RB1.1	+	-	-	x	<i>Staphylococcus sp.</i>
9	RB1.2	+	-	+	-	<i>Micrococcus luteus</i>
10	RB1.3	+	-	-	x	<i>Staphylococcus sp.</i>

Note: Isolate code represent bacterial isolates obtained from fecal samples of long-tailed macaques on Tinjil Island, including C72.2 (Cage 7 Sample 2 Isolate 2), C72.3 (Cage 7 Sample 2 Isolate 3), LB2.2 (Left Base camp Sample 2 Isolate 2), LB2.3 (Left Base camp Sample 2 Isolate 3), LB3.2 (Left Base camp Sample 3 Isolate 2), LB3.5 (Left Base camp Sample 3 Isolate 5), LB3.6 (Left Base camp Sample 3 Isolate 6), RB1.1 (Right Base camp Sample 1 Isolate 1), RB1.2 (Right Base camp Sample 1 Isolate 2), RB1.3 (Right Base camp Sample 1 Isolate 3). Types of biochemical tests: Ca (Catalase: (+) bubbles, (-) no bubbles); M (Mannitol: (+) yellow+bubbles in durham tube, (-) green); Y (Yellow pigment); G (Glucose: (+) yellow+bubbles in durham tube, (-) green). "x" indicates Not Tested

Table 2. Biochemical tests on bacterial isolates of Gram-positive bacilli

No	Isolate Code	Biochemical Test								Putative bacteria
		E	Af	Ca	S	VP	Cd	Mo	Na	
1	C72.1	+	x	+	+	-	x	x	-	<i>Bacillus sp.</i>
2	LB2.1	+	x	+	+	+	+	+	x	<i>Bacillus cereus</i>
3	LB2.4	-	-	+	+	x	x	x	x	<i>Corynebacterium kutscheri</i>
4	LB2.5	-	-	+	+	x	x	x	x	<i>Corynebacterium kutscheri</i>
5	LB3.1	-	-	+	+	x	x	x	x	<i>Corynebacterium kutscheri</i>
6	LB3.3	-	-	+	-	x	x	x	x	<i>Corynebacterium xerosis</i>
7	LB3.4	-	-	+	+	x	x	x	x	<i>Corynebacterium kutscheri</i>
8	RB1.4	-	-	+	-	x	x	x	x	<i>Corynebacterium xerosis</i>
9	RB1.5	+	x	+	+	+	+	+	x	<i>Bacillus cereus</i>

Note: The isolate code represents the code of bacterial isolates obtained from fecal samples of long-tailed macaques on Tinjil Island. C72.1 (Cage 7 Sample 2 Isolate 1), LB2.1 (Left Base camp Sample 2 Isolate 1), LB2.4 (Left Base camp Sample 2 Isolate 4), LB2.5 (Left Base camp Sample 2 Isolate 5), LB3.1 (Left Base camp Sample 3 Isolate 1), LB3.3 (Left Base camp Sample 3 Isolate 3), LB3.4 (Left Base camp Sample 3 Isolate 4), RB1.4 (Right Base camp Sample 1 Isolate 4), RB1.5 (Right Base camp Sample 1 Isolate 5). Type of biochemical tests E (Endospore stain: (+) endospore-forming, (-) non-endospore forming), Af (Acid-fast: (+) red cells, (-) blue cell, Ca (Catalase (+) bubbles, (-) no bubbles), S (Starch hydrolysis: (+) clear zone, (-) no clear zone), VP (Voges-Proskauer: (+) red, (-) orange/brown), Cd (Cell diameter: (+) Cd > 1 um, (-) Cd < 1 um), M (Motility test: (+) motile, (-) nonmotile), Na (NaCl 6.5%: (+) cloudy, (-) almost clear/same with negative control). "x" indicates Not Tested




Figure 4 Biochemical tests including Starch hydrolysis test and Hemolytic test. (Above) Results of Starch hydrolysis test for selected isolates: C72.1 (Cage 7 Sample 2 Isolate 1), LB2.1 (Left Base camp Sample 2 Isolate 1), LB2.4 (Left Base camp Sample 2 Isolate 4), LB2.5 (Left Base camp Sample 2 Isolate 5), LB3.1 (Left Base camp Sample 3 Isolate 1), LB3.3 (Left Base camp Sample 3 Isolate 3), LB3.4 (Left Base camp Sample 3 Isolate 4), RB1.4 (Right Base camp Sample 1 Isolate 4), RB1.5 (Right Base camp Sample 1 Isolate 5). (Below) Hemolytic activity of bacterial isolates from fecal samples of long-tailed macaques on Tinjil Island, categorized as (A)  $\alpha$ -hemolytic, (B)  $\beta$ -hemolytic, and (C)  $\gamma$ -hemolytic. B1 (C72.1/Cage 7 Sample 2 Isolate 1), C1 (C72.2/Cage 7 Sample 2 Isolate 2), C2 (C72.3/Cage 7 Sample 2 Isolate 3), B3 (LB2.1/Left Base camp Sample 2 Isolate 1), A2 (LB2.2/Left Base camp Sample 2 Isolate 2), A3 (LB2.3/Left Base camp Sample 2 Isolate 3), A4 (LB2.4/Left Base camp Sample 2 Isolate 4), A5 (LB2.5/Left Base camp Sample 2 Isolate 5), A6 (LB3.1/Left Base camp Sample 3 Isolate 1), C6 (LB3.2/Left Base camp Sample 3 Isolate 2), C7 (LB3.3/Left Base camp Sample 3 Isolate 3), A7 (LB3.4/Left Base camp Sample 3 Isolate 4), C8 (LB3.5/Left Base camp Sample 3 Isolate 5), A8 (LB3.6/Left Base camp Sample 3 Isolate 6), A1 (RB1.1/Right Base camp Sample 1 Isolate 1), C3 (RB1.2/Right Base camp Sample 1 Isolate 2), C4 (RB1.3/Right Base camp Sample 1 Isolate).

Based on the biochemical tests conducted on nine isolates of Gram-positive bacilli, two bacterial genera were identified: *Bacillus* and *Corynebacterium*, encompassing three putative species, namely *Bacillus cereus*, *Corynebacterium xerosis*, and *Corynebacterium kutscheri* (see Table 2, Figure 3, Supplementary Material Table S4, Table S5, Table S6). The endospore-forming bacterial isolates (LB2.1 and RB1.5) exhibited consistent results across all biochemical tests, indicating the presence of catalase and amylase enzymes (Figure 4), as well as acetoin production, as indicated by a positive VP test result.

Conversely, endospore-forming bacterial isolate C72.1 lacked the enzyme necessary for sugar breakdown into acetoin (negative VP test), differing from the other two endospore-forming isolates

(Supplementary Material Table S6). Isolate C72.1 was categorized as swollen cells due to the larger diameter of the spore compared to vegetative cells. Additionally, isolate C72.1 tested negative in the NaCl 0.65% test, indicating susceptibility to media conditions with NaCl 6.5%. Although two presumed bacterial species (*B. pantothenic* and *B. circulans*) were identified at this stage, the arabinose fermentation test could not be conducted, thus halting future identification.

Other bacterial isolates lacking endospore underwent shorter biochemical test pathways, including acid-fast staining, catalase test, and starch hydrolysis. Variations in starch hydrolysis test results distinguished six isolates into two different *Corynebacterium* species. Isolates forming clear zones (LB2.4, LB2.5, LB3.1, LB3.4) were identified



as *Corynebacterium kutscheri*, while those lacking clear zones (LB3.3 and RB1.4) were identified as *Corynebacterium xerosis* (Figure 4).

Hemolytic activity tests were conducted on all bacterial isolates, revealing various hemolytic types (Supplementary Material Table S6; Figure 4). Beta-hemolytic bacterial isolates formed clear zones around isolates, exemplified by isolates RB1.5, C72.1, and LB2.1 (Figure 4). Alpha-hemolytic bacteria were characterized by slightly darker streaks in the middle of quadrants, as observed in isolates LB3.1 and LB2.5 (Figure 4). Gamma hemolytic bacterial isolates exhibited no color change or alteration in the area surrounding the bacterial colony, as seen in LB3.5 and RB1.3. *Bacillus* isolates displayed beta-hemolytic activity, while other isolates, such as *Staphylococcus*, *Corynebacterium*, and *Micrococcus*, exhibited varying hemolytic patterns.

#### 4. Discussion

This study presents the findings from the isolation of bacteria from fecal samples of four semi-wild LTMs on Tinjil island. A total of 19 putative species of bacteria were identified, including three species of *Bacillus* spp., six of *Staphylococcus* spp. (belonging to the phylum Actinomycetota/Actinobacteria; Gao and Gupta 2012), four of *Micrococcus* spp., and six of *Corynebacterium* spp. (belong to phylum Bacilliota/Firmicutes; Ludwig *et al.* 2009). Typically, the predominant bacterial phyla in the gut of NHPs include Firmicutes, Bacteroidetes, Proteobacteria, and others (Gogarten *et al.* 2018). Notably, Firmicutes, Bacteroidetes, Proteobacteria, and Tenericutes are highly common in LTMs (Grant *et al.* 2019; Nagpal *et al.* 2018; Sawaswong *et al.* 2021). These findings align with previous studies employing similar methodologies in baboons (Lugano *et al.* 2018). Subsequently, based on the

intestinal oxygen concentration and bacterial types, it can be inferred that bacterial genera such as *Bacillus* (Toerien 1967), *Staphylococcus* and *Micrococcus* (Kloos and Musselwhite 1975), and *Corynebacterium* (Nishimura *et al.* 2011) can survive in the intestine under suitable oxygen concentrations conducive to the growth of these types of bacteria.

The *Bacillus* species identified in our study encompassed two putative species: *Bacillus cereus* and *Bacillus* spp. *Bacillus cereus* is known as an emetic and enterotoxin-producing bacterium capable of causing diarrhea (Benedict *et al.* 1993) and has been characterized as a beta-hemolytic (Dabiré *et al.* 2022). *Bacillus* spp. is found in various habitats, including the gut of various insects and animals (Hong *et al.* 2009), and generally exhibits the ability to hydrolyze starch and protein in anaerobic digesters (Toerien 1967). Consequently, *Bacillus* spp. are used in numerous medical, pharmaceutical, agricultural, and industrial processes due to their broad range of physiological characteristics, including the production of enzymes such as amylase (Luang-In *et al.* 2019) and antibiotics (Beneduzi *et al.* 2012). While certain species of *Bacillus*, notably *Bacillus cereus*, are known to be occasional pathogens of humans and livestock, most *Bacillus* spp. are harmless saprophytes (Turnbull *et al.* 2002). In the case of semi-wild LTMs, their diet may consist of plants/fruits, insects, and soil. Spores from *Bacillus cereus* are known to proliferate in the intestines of insects (Margulis *et al.* 1998), which may then be ingested by LTMs, ultimately resulting in the colonization of these bacteria in their intestines. Although one of the putative bacterial species of *Bacillus* identified in this study was classified as pathogenic, there is a possibility that *Bacillus* spp. represent beneficial species of *Bacillus*, such as *Bacillus coagulans*. This speculation arises because all identified bacterial species are still categorized as

putative. Furthermore, *Bacillus coagulans* can serve as a probiotic, alleviating clinical symptoms such as bloating, vomiting, diarrhea, and abdominal pain in patients with irritable bowel syndrome (Majeed *et al.* 2016).

The only putative species identified within the *Staphylococcus* genera were *Staphylococcus aureus* and *Staphylococcus* spp. *Staphylococcus* is known to inhabit various environments, including the air (Solberg 2000), animal skins (squirrel, monkey, and sheep) (Kloos *et al.* 1976), and human skin (Raineri *et al.* 2022). *Staphylococcus aureus* is the causative agent of food poisoning, one of the most common foodborne illnesses that can induce nausea (vomiting) and diarrhea (Abril *et al.* 2020). However, the presence of some putative bacterial species in these semi-wild LTM samples does not necessarily imply that the conditions of the LTMs on Tinjil Island are unhealthy or that they are suffering from diarrhea. The results obtained in this study do not permit a full assessment of the complete condition of the gut microbiota of the LTMs on Tinjil with respect to their health, including whether they are experiencing dysbiosis. Nevertheless, direct field observations indicate that the LTMs did not suffer from diarrhea, as evidenced by the stool condition observed during fecal sampling, which exhibited a normal texture (soft and well-formed in consistency).

In addition to *Staphylococcus*, it appears that *Micrococcus* is also a genus of microbes commonly found on the skin of animals, in the air, within the inner tissues of plants, in soil, and among several fish species (Lee *et al.* 2022). The only putative species identified in this study, namely *Micrococcus luteus*, is categorized as human-commensal and non-pathogenic bacterial species (Albertson *et al.* 1978). This study also identified other putative bacterial species, namely *Corynebacterium xerosis* and *Corynebacterium*

*kutscheri*. *Corynebacterium xerosis* is a commensal organism typically present on the skin and mucous membranes of humans and animals (Vela *et al.* 2006), while *Corynebacterium kutscheri* has been described as a commensal bacterium in mice, rats, and voles (Holmes and Korman 2007). The identified putative bacterial species obtained in this study exhibit a range of characteristics, being either non-pathogenic, pathogenic, or commensal to their host. However, only a few (five putative species) could be identified up to the species level in this study.

The limited number of putative bacterial species obtained in this study can be attributed to the small number of bacterial isolates per sample. Among the four fecal samples analyzed, C72 exhibited the lowest bacterial isolates compared to LB2, LB3, and RB1. This scarcity of bacterial isolates in C72 occurred because there were only very few distinct bacterial colonies when isolating bacteria from a solid medium. Identifying certain bacteria enables the discrimination between different bacterial species based on their morphology, arrangement, and consequent colony patterns (Badieyan *et al.* 2018).

The incubation technique used in this study also likely played a role in affecting the lower diversity of isolated bacteria in this study. All procedures, including sampling collection, preparation, and isolation, were conducted aerobically without implementing anaerobic conditions. Consequently, only aerobic or facultative anaerobic bacteria could thrive, while obligate anaerobic bacteria could not. Ideally, bacterial isolation should encompass both aerobic and anaerobic conditions to facilitate the growth of a wider range of bacteria. Moreover, not only should fecal sample collection and storage be performed anaerobically in the laboratory, but it is imperative due to the dominant presence of anaerobic bacteria, including Bacteroidetes, Firmicutes, and

Proteobacteria, comprising over 95% of rectal microbiota in rhesus and LTM macaques (Cui *et al.* 2019). Therefore, exposure to oxygen should be carefully considered as an influential factor.

The isolation of cultivable fecal microbiota from semi-wild LTMs on Tinjil Island has provided a basic overview and preliminary baseline data of gut bacterial diversity through culture-dependent methods. However, studies examining intestinal bacteria diversity through morphological and biochemical analysis have inherent limitations compared to molecular analysis. Non-molecular methods can only culture bacteria capable of growth in vitro conditions, thus restricting the range of detectable bacteria (Hayashi *et al.* 2022). Therefore, molecular approaches, such as metagenomic analysis, are essential for a comprehensive understanding of gut microbiota profiles. Molecular analysis offers increased sensitivity and accuracy in identifying bacterial species compared to culture-dependent methods (McKenna *et al.* 2008; Rhoads *et al.* 2012).

Nevertheless, intestinal bacteria play a crucial role as key regulators of digestion throughout the gastrointestinal tract (Bornbusch *et al.* 2023; Rinninella *et al.* 2019; Turnbaugh and Gordon 2009). Various factors influence the diversity and composition of gut microbiota, with diet being the most critical determinant in NHPs (Gogarten *et al.* 2018; Lee *et al.* 2023; Nagpal *et al.* 2018). Our findings hold practical implications for management facilities, particularly in cage-breeding facilities, by guiding ex-situ food management protocols. Consequently, data on intestinal microbiota profiles can inform strategies for regulating nutritional intake for captive or wild primates. This could involve incorporating higher-quality natural fiber-rich foods or probiotics into the diet of captive LTM (Muhammad

*et al.* 2023 ; Tian *et al.* 2022). Therefore, studies on gut microbiota profiles in NHPs, whether in wild or captive settings, are crucial for informing primate welfare management practices and will be invaluable for future management efforts.

## Acknowledgments

The authors express their gratitude to the Young Lecturer Research Grant (*Penelitian Dosen Muda*) 2023, IPB University, for providing financial support for this study. Additionally, we extend our appreciation to the Primate Research Center, IPB University, and the Laboratory of Microbiology Department of Biology, IPB University for granting permission to conduct this study. RCK's effort was supported in part by the Office of Research Infrastructure Programs (ORIP) of the National Institutes of Health through grant number P51OD010425 to the Washington National Primate Research Center, USA.

## References

- Abril, A.G., Villa. T.G., Barros-Velázquez, J., Cañas, B., Sánchez-Pérez, A., Calo-Mata, P., Carrera, M. 2020. *Staphylococcus aureus* exotoxins and their detection in the dairy industry and mastitis. *Toxins (Basel)*, 12(9), 1–18. <https://doi.org/10.3390/toxins12090537>.
- Albertson, D., Natsios, G.A., Gleckman, R. 1978. Septic shock with *Micrococcus luteus*. *Arch. Intern. Med.*, 138(3), 487-495. <https://doi.org/10.1001/archinte.1978.03630270093032>.
- Badieyan, S., Dilmaghani-Marand, A., Hajipour, M.J., Ameri, A., Razzaghi, M.R., Rafii-Tabar, H., Mahmoudi, M., Sasanpour, P. 2018. Detection and discrimination of bacterial colonies with mueller matrix imaging. *Sci. Rep.*, 8(1), 1–11. <https://doi.org/10.1038/s41598-018-29059-5>.

- Benedict, R.C., Partridge, T., Wells, D., Buchanan, R.L. 1993. *Bacillus cereus*: Aerobic growth kinetics. *J. Food Prot.*, 56(3), 211–214. <https://doi.org/10.4315/0362-028X-56.3.211>.
- Beneduzi, A., Ambrosini, A., Passaglia, L.M.P. 2012. Plant growth-promoting rhizobacteria (PGPR): Their potential as antagonists and biocontrol agents[review]. *Genet. Mol. Biol.*, 4, 1044–1051. <https://doi.org/10.1590/s1415-47572012000600020>.
- Boonkusol, D., Thongyuan, S., Jangsuwan, N., Sanyathitiseree, P. 2020. Antimicrobial resistance profiles in bacterial species isolated from fecal samples of free-ranging long-tailed macaques (*Macaca fascicularis*) living in Lopburi Old Town, Thailand. *Vet. World.*, 13(7), 1397–1403. <https://doi.org/10.14202/vetworld.2020.1397-1403>.
- Bornbusch, S.L., Muletz-Wolz, C.R., Lopez-Bondarchuck, E., Maslanka, M.T., Kendrick, E.L. 2023. Gut microbiomes of captive primates show phylosymbiosis, respond to dietary sugar reduction, and select for host-specific dietary microbes. *FEMS Microbiol. Ecol.*, 99, 1–11. <https://doi.org/10.1093/femsec/fiad069>.
- Bruins, M.J., Juffer, P., Wolfhagen, M.J.H.M., Ruijs, G.J.H.M. 2007. Salt tolerance of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*. *J. Clin. Microbiol.*, 45(2), 682–683. <https://doi.org/10.1128/jcm.02417-06>.
- Buxton, R. 2016. Blood agar plates and hemolysis protocols. American Society for Microbiology. <https://asm.org/protocols/blood-agar-plates-and-hemolysis-protocols>. [Date accessed: 11 December 2023].
- Carabotti, M., Scirocco, A., Maselli, M.A., Severi, C. 2015. The gut-brain axis: Interactions between enteric microbiota, central and enteric nervous systems. *Ann. Gastroenterol.*, 28(2), 203–209.
- Carbia, C., Bastiaanssen, T.F.S., Iannone, L.F., García-Cabrerizo, R., Boscaini, S., Berding, K., Strain, C.R., Clarke, G., Stanton, C., Dinan, T.G., Cryan, J.F. 2023. The Microbiome-Gut-Brain axis regulates social cognition & craving in young binge drinkers. *eBioMedicine*, 89, 104442. <https://doi.org/10.1016/j.ebiom.2023.104442>.
- Chen, T., Long, W., Zhang, C., Liu, S., Zhao, L., Hamaker, B.R. 2017. Fibre-utilising capacity varies in *Prevotella*- versus *Bacteroides*-dominated gut microbiota. *Sci. Rep.*, 7(1), 2594. <https://doi.org/10.1038/s41598-017-02995-4>.
- Cui, Y.F., Wang, F.J., Yu, L., Ye, H.H., Yang, G.B. 2019. Metagenomic comparison of the rectal microbiota between rhesus macaques (*Macaca mulatta*) and cynomolgus macaques (*Macaca fascicularis*). *Zool. Res.*, 40(2), 89–93. <https://doi.org/10.24272%2Fj.issn.2095-8137.2018.061>.
- Dabiré, Y., Somda, N.S., Somda, M.K., Mogmenga, I., Traoré, A.K., Ezeogu, L.I., Traoré, A.S., Ugwuanyi, J.O., Dicko, M.H. 2022. Molecular identification and safety assessment of *Bacillus* strains isolated from Burkinabe traditional condiment “soumbala.” *Ann. Microbiol.*, 72(1), 10–20. <https://doi.org/10.1186/s13213-022-01664-w>.
- Frankel, J.S., Mallott, E.K., Hopper, L.M., Ross, S.R., Amato, K.R. 2019. The effect of captivity on the primate gut microbiome varies with host dietary niche. *Am. J. Primatol.*, 81(12), e23061. <https://doi.org/10.1002/ajp.23061>.



- Gao, B., Gupta, R.S. 2012. Phylogenetic framework and molecular signatures for the main clades of the Phylum Actinobacteria. *Microbiol. Mol. Biol.*, 76(1), 66–112. <https://doi.org/10.1128%2FMMBR.05011-11>.
- Gogarten, J.F., Davies, T.J., Benjamino, J., Gogarten, J.P., Graf, J., Mielke, A., Mundry, R., Nelson, M.C., Wittig, R.M., Leendertz, F.H., Calvifnac-Spencer, S. 2018. Factors influencing bacterial microbiome composition in a wild non-human primate community in Taï National Park, Côte d'Ivoire. *ISME.*, 12(10), 2559–2574. <https://doi.org/10.1038%2Fs41396-018-0166-1>.
- Grant, E.T., Kyes, R.C., Kyes, P., Trinh, P., Ramirez, V., Tanee, T., Pinlaor, P., Dangtakot, R., Rabinowitz, P.M. 2019. Correction: Faecal microbiota dysbiosis in macaques and humans within a shared environment. *PLoS One.*, 14(5), e0210679. <https://doi.org/10.1371/journal.pone.0210679>.
- Hansen, M.F., Ang, A., Trinh, T., Sy, E., Paramasiwam, S., Ahmed, T., Dimalibot, J., Jones-Engel, L., Ruppert, N., Griffioen, C., Gray, R., Phiapalath, P., Doak, N., Kite, S., Nijman, V., Fuentes, A., Gumert, M.D. 2022. *Macaca fascicularis*. The IUCN Red List of Threatened Species 2022: e.T12551A199563077. 8235. <https://www.iucnredlist.org/fr/species/12551/221666136>. [Date accessed: 15 December 2022]
- Hanson, A. 2008. Oxidative-fermentation test. American Society for Microbiology. <https://asm.org/protocols/oxidative-fermentative-test-protocol>. [Date accessed: 15 December 2022].
- Hasanah, R., Maulana, V.S., Iskandar, E. 2022. Sexual behaviour of long-tailed macaques (*Macaca fascicularis*) in semi-natural captivity, Tinjil Island, Indonesia. *InaJP.*, 1(1), 1–9. <https://doi.org/10.29244/primatology.1.01.1-9>.
- Hayashi, T., Ichikawa, M., Konishi, I. 2022. Spontaneous myocarditis in mice predisposed to autoimmune disease: Including vaccination-induced onset. *Biomedicines*, 10(6), 1443. <https://doi.org/10.1101/2022.03.14.484354>.
- Hilborn, R., Smith, D.R. 2024. Is the long-tailed macaque at risk of extinction?. *Am. J. Primatol.*, 86(4), e23590. <https://doi.org/10.1002/ajp.23590>.
- Holmes, N.E., Korman, T.M. 2007. *Corynebacterium kutscheri* infection of skin and soft tissue following rat bite. *J. Clin. Microbiol.*, 45(10), 3468–3469. <https://doi.org/10.1128/jcm.00607-07>.
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T., Williams, S.T. 1994. *Bergey's Manual of Determinative Bacteriology*. Maryland: Lippincott Williams & Wilkins.
- Hong, H.A., To, E., Fakhry, S., Baccigalupi, L., Ricca, E., Cutting, S.M. 2009. Defining the natural habitat of *Bacillus* spore-formers. *Res. Microbiol.*, 160(6), 375–379. <https://doi.org/10.1016/j.resmic.2009.06.006>.
- Hussey, M.A., Zayaitz, A. 2008. Acid-fast stain protocols. American Society for Microbiology. <https://asm.org/ASM/media/Protocol-Images/Acid-Fast-Stain-Protocols.pdf?ext=.pdf>. [Date accessed: 9 December 2022].
- Hussey, M. A. and Zayaitz, A. 2007. Endospore stain protocol, American Society for Microbiology. <https://www.asmscience.org/content/education/protocol/protocol.3112>. [Date accessed: 10 Desember 2022].
- Kloos, W.E., Musselwhite, M.S. 1975. Distribution and persistence of *Staphylococcus* and *Micrococcus* species and other aerobic bacteria on human skin. *Appl. Microbiol.*, 30(3), 381–395. <https://doi.org/10.1128/am.30.3.381-395.1975>.

- Kloos, W.E., Zimmerman, R.J., Smith, R.F. 1976. Preliminary studies on the characterization and distribution of *Staphylococcus* and *Micrococcus* species on animal skin. *AEM*, 31(1), 53–59. <https://doi.org/10.1128/aem.31.1.53-59.1976>.
- Koo, B.S., Baek, S.H., Kim, G., Hwang, E., Oh, H., Son, Y., Lim, K.S., Kang, P., Lee, H.Y., Jeong, K., Kim, Y., Villinger, F., Hong, J. 2020. Idiopathic chronic diarrhoea associated with dysbiosis in a captive cynomolgus macaque (*Macaca fascicularis*). *J. Med. Primatol.*, 49(1), 56–59. <https://doi.org/10.1111/jmp.12447>.
- Kyes, R.C. 1993. Survey of the long-tailed macaques introduced onto Tinjil Island, Indonesia. *Am. J. Primatol.*, 31(1), 77–83. <https://doi.org/10.1002/ajp.1350310108>.
- Lee, A.Y., Chen, C.H., Liou, J.S., Lin, Y.C., Hamada, M., Wang, Y.T., Peng, L.L., Chang, S.C., Chen, C.C., Lin, C.F., Huang, L., Huang, CH. 2022. *Micrococcus porci* sp. nov., isolated from feces of black pig (*Sus scrofa*). *Life (Basel)*, 12(11), 1749. <https://doi.org/10.3390/life12111749>.
- Lee, W., Oi, T., Kondo, T., Uno, T., Seki, K., Smiada, M., Tsuji, Y., Langgeng, A., macIntosh, A., Suzuki, K., Yamada, K., Onishi, K., Ueno, M., Kubo, K., Hanya, G. 2023. Diet-related factors strongly shaped the gut microbiota of Japanese macaques. *Am. J. Primatol.*, 85(12), e23555. <https://doi.org/10.1002/ajp.23555>.
- Li, H.Z., Li, N., Wang, J.J., Li, H., Huang, X., Guo, L., Zheng, H.W., He, Z.L., Zhao, Y., Yang, Z.N., Fan, H.T., Chu, M.M., Yang, J.X., Wu, Q.W., Liu, L.D. 2020. Dysbiosis of gut microbiome affecting small intestine morphology and immune balance: Arhesus macaque model. *Zool. Res.*, 41(1), 20–31. <https://doi.org/10.24272/j.issn.2095-8137.2020.004>.
- Li, X., Santos, R., Bernal, J.E., Li, D.D., Hargaden, M., Khan, N.K. 2023. Biology and postnatal development of organ systems of cynomolgus monkeys (*Macaca fascicularis*). *J. Med. Primatol.*, 52(1), 64–78. <https://doi.org/10.1111/jmp.12622>.
- Liedigk, R., Kolleck, J., Böker, K.O., Meijaard, E., Md-Zain, B.M., Abdul-Latiff, M.A.B., Ampeng, A., Lakim, M., Abdul-Patah, P., Tosi, A.J., Brameier, M., Zinner, D., Ross, C. 2015. Mitogenomic phylogeny of the common long-tailed macaque (*Macaca fascicularis fascicularis*). *BMC Genomics*, 16(1), 222. <https://doi.org/10.1186/s12864-015-1437-0>.
- Luang-In, V., Yotchaisarn, M., Saengha, W., Udomwong, P., Deeseenthum, S., Maneewan, K. 2019. Isolation and identification of amylase-producing bacteria from soil in Nasinuan community forest, Maha Sarakham, Thailand. *BPJ*, 12(3), 1061–1068. <http://dx.doi.org/10.13005/bpj/1735>.
- Ludwig, W., Schleifer, K.H., Whitman, W.B. 2009. Revised road map to the phylum Firmicutes. In: *De Vos, P., et al. Bergey's Manual® of Systematic Bacteriology*. New York, USA: Springer.
- Lugano, S. D., Nyerere, K. A., Kariuki, W. K., Samuel, K., Joseph, K., & Apondi, O. J. 2018. Gastrointestinal microbial flora in wild and captive olive baboons (*Papio anubis*). *Am. J. Infect. Dis.*, 6(1), 30-37. <http://dx.doi.org/10.12691/ajidm-6-1-5>.
- Majeed, M., Nagabhushanam, K., Natarajan, S., Sivakumar, A., Ali, F., Pande, A., Majeed, S., Karri, S.K. 2016. *Bacillus coagulans* MTCC 5856 supplementation in the management of diarrhoea predominant Irritable Bowel Syndrome: A double blind randomised placebo controlled pilot clinical study. *Nutr. J.*, 15(1), 21–30. <https://doi.org/10.1186/s12937-016-0140-6>.

- Margulis, L., Jorgensen, J.Z., Dolan, S., Kolchinsky, R., Rainey, F.A., Lo, S.C. 1998. The arthromitus stage of *Bacillus cereus*: Intestinal symbionts of animals. *Proc. Natl. Acad. Sci. U S A.*, 95(3), 1236–1241. <https://doi.org/10.1073/pnas.95.3.1236>.
- Martin, A.M., Sun, E.W., Rogers, G.B., Keating, D.J. 2019. The influence of the gut microbiome on host metabolism through the regulation of gut hormone release. *Front. Physiol.*, 10(3), 1–11. <https://doi.org/10.3389%2Ffphys.2019.00428>.
- McDevitt, S. 2009. *Methylred and voges-proskauer test protocols*. American Society for Microbiology. <https://asm.org/protocols/methyl-red-and-voges-proskauer-test-protocols>. [Date accessed: 10 December 2022].
- McKenna, P., Hoffmann, C., Minkah, N., Aye, P.P., Lackner, A., Liu, Z., Lozupone, C.A., Hamady, M., Knight, R., Bushman, F.D. 2008. The macaque gut microbiome in health, lentiviral infection, and chronic enterocolitis. *PLoS Pathog.*, 4(2), e20. <https://doi.org/10.1371%2Fjournal.ppat.0040020>.
- Muegge, B.D., Kuczynski, J., Knights, D., Clemente, J.C., González, A., Fontana, L., Henrissat, B., Knight, R., Gordon, J.I. 2011. Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science*, 332(6032), 970–974. <https://doi.org/10.1126/science.1198719>.
- Muhammad, R., Klomkliev, P., Chanchaem, P., Sawaswong, V., Kaikaew, T., Payungporn, S., Malaivijitnond, S. 2023. Comparative analysis of gut microbiota between common (*Macaca fascicularis fascicularis*) and Burmese (*M. f. aurea*) long-tailed macaques in different habitats. *Sci. Rep.*, 13(1), 14950. <https://doi.org/10.1038%2Fs41598-023-42220-z>.
- Nagpal, R., Mainali, R., Ahmadi, S., Wang, S., Singh, R., Kavanagh, K., Kitzman, D.W., Kushugulova, A., Marotta, F., Yadav, H. 2018. Gut microbiome and aging: Physiological and mechanistic insights. *Nutr. Healthy Aging.*, 4(4), 267–285. <https://doi.org/10.3233/nha-170030>.
- Nishimura, T., Teramoto, H., Inui, M., Yukawa, H. 2011. Gene expression profiling of *Corynebacterium glutamicum* during anaerobic nitrate respiration: Induction of the SOS response for cell survival. *J. Bacteriol.*, 193(6), 1327–1333. <http://dx.doi.org/10.1128/JB.01453-10>.
- Pamungkas, J., Sajuthi, D., Lelana P.A., Iskandriati, D., Joeniman, B., Kyes, R.C., Knitter, G.H., Wantanabe, R.A. 1994. Tinjil island, a natural habitat breeding facility of simian retrovirusfree *Macaca fascicularis*. *Am. J. Primatol.*, 34(1), 81–84. <https://doi.org/10.1002/ajp.1350340113>.
- Perwitasari-Farajallah, D., Iskandar, E., Sawitri, H.I., Abimanyu, T.L., Maulana, V.S., Rachmawati, A.D., Purnama, I., Darusman, H.S. 2023a. Population estimate of long-tailed macaques (*Macaca fascicularis*) on Tinjil Island. *HAYATI J Biosci.*, 30(2), 193–197. <https://doi.org/10.4308/hjb.30.2.193-197>.

- Perwitasari-Farajallah, D., Iskandar, E., Sawitri, H.I., Abimanyu, T.L., Maulana, V.S., Rachmawati, A.D., Purnama, I., Darusman, H.S. 2023a. Population estimate of long-tailed macaques (*Macaca fascicularis*) on Tinjil Island. *HAYATI J Biosci.*, 30(2), 193–197. <https://doi.org/10.4308/hjb.30.2.193-197>.
- Pokusaeva, K., Johnson, C., Luk, B., Uribe, G., Fu, Y., Oezguen, N., Matsunami, R.K., Lugo, M., Major, A., Mori-Akiyama, Y., Hollister, E.B., Dann, S.M., Shi, X.Z., Engler, D.A., Savidge, T., Versalovic, J. 2017. GABA-producing *Bifidobacterium dentium* modulates visceral sensitivity in the intestine. *Neurogastroenterol. Motil.*, 29(1), e12904. <https://doi.org/10.1111/nmo.12904>.
- Prats, J., Garcia-Armisen, T., Larrea, J., Servais, P. 2008. Comparison of culture-based methods to enumerate *Escherichia coli* in tropical and temperate freshwaters. *Lett. Appl. Microbiol.*, 46(2), 243-248. <https://doi.org/10.1111/j.1472-765x.2007.02292.x>.
- Raineri, E.J.M., Maaß, S., Wang, M., Brushett, S., Medina, L.M.P., Escandell, N.S., Altulea, D., Raangs, E., de Jong, A., Murguia, E.V., Feil, E.J., Friedrich, A.W., Buist, G., Becher, D., Garcia-Cobos, S., Couto, N., van Dijk, J.M. 2022. *Staphylococcus aureus* populations from the gut and the blood are not distinguished by virulence traits a critical role of host barrier integrity. *Microbiome*, 10(1), 239-262. <https://doi.org/10.1186/s40168-022-01419-4>.
- Reiner, K. 2010. *Catalase test protocol*. American Society for Microbiology. <https://asm.org/protocols/catalase-test-protocol>. [Date accessed: 10 December 2023].
- Rhoads, D.D., Wolcott, R.D., Sun, Y., Dowd, S.E. 2012. Comparison of culture and molecular identification of bacteria in chronic wounds. *Int. J. Mol. Sci.*, 13(3), 2535–2550. <https://doi.org/10.3390%2Fijms13032535>.
- Rinninella, E., Raoul, P., Cintoni, M., Franceschi, F., Miggiano, G.A.D., Gasbarrini, A., Mele, M.C. 2019. What is the healthy gut microbiota composition? A changing ecosystem across age, environment, diet, and diseases. *Microorganisms*, 7(1), 14-35. <https://doi.org/10.3390%2Fmicroorganisms7010014>.
- Sawaswong, V., Chanchaem, P., Kemthong, T., Warit, S., Chaiprasert, A., Malaivijitnond, S., Payungporn, S. 2023. Alteration of gut microbiota in wild - borne long - tailed macaques after 1 - year being housed in hygienic captivity. *Sci. Rep.*, 15(1), 5842. <https://doi.org/10.1038/s41598-023-33163-6>.
- Sawaswong, V., Praianantathavorn, K., Chanchaem, P., Khamwut, A., Kemthong, T., Hamada, Y., Malaivijitnond, S., Payungporn, S. 2021. Comparative analysis of oral-gut microbiota between captive and wild long-tailed macaque in Thailand. *Sci. Rep.*, 11(1), 14280. <https://doi.org/10.1038/s41598-021-93779-4>.
- Smith, A., Hussey, M. 2020. *Gram stain protocols*. American Society for Microbiology. <https://asm.org/protocols/gram-stain-protocols>. [Date accessed: 10 December 2022].
- Solberg, C.O. 2000. Spread of *Staphylococcus aureus* in hospitals: Causes and prevention. *Scand. J. Infect. Dis.*, 32(6), 587–595. <https://doi.org/10.1080/003655400459478>.



- Stevison, L.S., Kohn, M.H. 2008. Determining genetic background in captive stocks of cynomolgus macaques (*Macaca fascicularis*). *J. Med. Primatol.*, 37(6), 311–317. <https://doi.org/10.1111/j.1600-0684.2008.00292.x>.
- Takenaka, A., Matsumoto, Y., Nagaya, A., Watanabe, K., Goto, S., Suryobroto, B., Takenaka, O. 2000. Plasma cholesterol levels in free-ranging macaques compared with captive macaques and humans. *Primates*, 41(3), 299–309. <https://doi.org/10.1007/bf02557599>.
- Tian, P., Gao, J., Liang, L., Cui, B., Hu, Q., Zhou, W., Li, B., Liu, Y., Chen, T., Rao, J., Wei, H. 2022. Faecal microbiota transplantation could improve chronic diarrhea in cynomolgus monkey by alleviating inflammation and modulating gut microbiota. *Biomedicines*, 10(12), 3016. <https://doi.org/10.3390/biomedicines10123016>.
- Toerien, D.F. 1967. Enrichment culture studies on aerobic and facultative anaerobic bacteria found in anaerobic digesters. *Water Res.*, 1(2), 147-155. [https://doi.org/10.1016/0043-1354\(67\)90081-4](https://doi.org/10.1016/0043-1354(67)90081-4). 155.
- Turnbaugh, P.J., Gordon, J.I. 2009. The core gut microbiome, energy balance and obesity. *Physiol. J.*, 587(17), 4153–4158. <https://doi.org/10.1113/jphysiol.2009.174136>.
- Turnbull, P.C.B., Quinn, C.P., Henderson, I. 2002. *Bacillus anthracis* and other *Bacillus* species. In Sussman, M. (Ed.), *Molecular Medical Microbiology* (pp. 2011-2031). Massachusetts, USA: Academic Press.
- Vela, A.I., Gracia, E., Fernández, A., Domínguez, L., Fernández-Garayzábal, J.F. 2006. Isolation of *Corynebacterium xerosis* from animal clinical specimens. *J. Clin. Microbiol.*, 44(6), 2242–2243. <https://doi.org/10.1128/jcm.02473-05>.
- Wills, M.O., Shields-Cutler, R.R., Brunmeier, E., Weissenborn, M., Murphy, T., Knights, D., Johnson, T.J., Clayton, J.B. 2022. Host species and captivity distinguish the microbiome compositions of a diverse zoo-resident nonhuman primate population. *Diversity*, 14(9), 715-730. <https://doi.org/10.3390/d14090715>.
- Wu, J., Chai, T., Zhang, H., Huang, Y., Perry, S.W., Li, Y., Duan, J., Tan, X., Hu, X., Liu, Y., Pu, J., Wang, H., Song, J., Jin, X., Ji, P., Zheng, P., Xie, P. 2022. Changes in gut viral and bacterial species correlate with altered 1,2-diacylglyceride levels and structure in the prefrontal cortex in a depression-like non-human primate model. *Transl. Psychiatry*, 12(1), 74-84. <https://doi.org/10.1038/s41398-022-01836-x>.
- Yano, J.M., Yu, K., Donaldson, G.P., Shastri, G.G., Ann, P., Ma, L., Nagler, C.R., Ismagilov, R.F., Mazmanian, S.K., Hsiao, E.Y. 2015. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*, 161(2), 264–276. <https://doi.org/10.1016/j.cell.2015.02.047>.
- Ying, C., Siao, Y., Chen, W., Chen, Y-T., Chen, S., Chen, Y-L., Hsu, J. 2022. Host species and habitats shape the bacterial community of gut microbiota of three non-human primates: Siamangs, white-handed gibbons, and Bornean orangutans. *Front. Microbiol.*, 13(1), 920190. <https://doi.org/10.3389/fmicb.2022.920190>.
- Zuberer, D.A. 1994. Recovery and enumeration of viable bacteria. In Weaver, R.W., Angle, S., Bottomley, P., Bezdicek, D., Smith, S., Tabatabai, A., Wollum, A. (Eds). *Methods soil analysis: Part 2 microbiological and biochemical properties* (pp.119–144). New York, USA: John-Wiley & Sons.